



TOPSCIENCE 众山



L-Ascorbic Acid 2-Glucoside

Material Submission Code:
004104-00507-6011

SHANDONG TOPSCIENCE BIOTECH CO., LTD.

Address: #98 Lanshan West Road, Lanshan District, Rizhao, Shandong, China

Tel: +86-532-80821298

E-mail: sales@topscience.cn

Website: www.topscience.cn

QINGDAO TRONGEN BIOTECH CO., LTD.

Address: #330 Songling Road, Laoshan District, Qingdao, Shandong, China

Tel: +86-532-80821298

E-mail: sales@trongenbio.com

Website: www.topscience.cn




INCI NAME: ASCORBYL GLUCOSIDE

MOLECULAR WEIGHT: 338.26

MOLECULAR FORMULA: C₁₂H₁₈O₁₁

RECOMMENDED DOSAGE: 0.1%~2%



PRODUCT FEATURE



L-Ascorbic Acid 2-Glucoside is a kind of natural Vitamin C containing stable glucose stabilizers.




L-Ascorbic Acid 2-Glucoside has good stability and solubility in solution. It can essentially inhibit discoloration and degradation of Vitamin C in cosmetics while retaining all the biological activity of Vitamin C.



L-Ascorbic Acid 2-Glucoside can desalinate pigment and reduce the pigmentation of senile spots and freckles; Brighten skin tone, delaying aging, protecting the skin in sunscreen products.


EFFICACY TEST

01 L-Ascorbic Acid 2-Glucoside Can Stimulate Collagen Synthesis




The results showed that compared with the control group, the L-Ascorbic Acid 2-Glucoside group had a greater stimulating effect on collagen synthesis, which could stimulate collagen synthesis and delay aging.

02 L-Ascorbic Acid 2-Glucoside Inhibits Melanin Synthesis



The results showed that both L-Ascorbic Acid 2-Glucoside and Vitamin C have the effect of inhibiting melanin synthesis, and the inhibitory effect of L-Ascorbic Acid 2-Glucoside is better than that of Vitamin C.



The results showed that both L-Ascorbic Acid 2-Glucoside and Vitamin C could inhibit melanin synthesis. And the inhibitory effect of L-Ascorbic Acid 2-Glucoside is superior to that of Vitamin C.

APPLICATION

Mainly applied in cosmetics, such as lotions, creams, and sunscreens.

PACKAGE AND STORAGE

1. 1kg/bag or 5kg/bag, pharmaceutical low density polyethylene bag with outer aluminum foil bag.
2. In an airtight container, protected from light and humidity, 2-8°C.